

BAHAGIAN KEJURURAWATAN  
KEMENTERIAN KESIHATAN MALAYSIA

**HUMPTY DUMPTY FALL RISK ASSESSMENT SCALE CHART(PAEDIATRIC)**

| Name:  |       | Age:  |       |       |       |       |       |       |
|--|-------|-------|-------|-------|-------|-------|-------|-------|
| RN No:   |       | Ward: |       |       |       |       |       |       |
| PARAMETER  | DATE  |       |       |       |       |       |       |       |
|  | TIME  |       |       |       |       |       |       |       |
| Age  | SCORE | SCORE | SCORE | SCORE | SCORE | SCORE | SCORE | SCORE |
| <3 years old   | 4     |       |       |       |       |       |       |       |
| 3 years to < 7 years old   | 3     |       |       |       |       |       |       |       |
| 7 years to , 13 years old  | 2     |       |       |       |       |       |       |       |
| 13 years +   | 1     |       |       |       |       |       |       |       |
| Gender   |       |       |       |       |       |       |       |       |
| Male   | 2     |       |       |       |       |       |       |       |
| Female   | 1     |       |       |       |       |       |       |       |
| Diagnosis  |       |       |       |       |       |       |       |       |
| Neurological Diagnosis   | 4     |       |       |       |       |       |       |       |
| Alteration in oxygenation eg<br>Respiratory Diagnosis,<br>Dehydration, Anaemia,<br>Syncope/Dizziness etc.                      | 3     |       |       |       |       |       |       |       |
| Psychological/Behavioral<br>Disorders  | 2     |       |       |       |       |       |       |       |
| Other Diagnosis  | 1     |       |       |       |       |       |       |       |
| Cognitive Impairment   |       |       |       |       |       |       |       |       |
| Not aware of limitations   | 3     |       |       |       |       |       |       |       |
| Forgets limitations  | 2     |       |       |       |       |       |       |       |
| Orientated to own ability  | 1     |       |       |       |       |       |       |       |
| Environmental Factors  |       |       |       |       |       |       |       |       |
| History of falls Infant -<br>Toddler placed in bed   | 4     |       |       |       |       |       |       |       |
| Patient uses assistive devices   | 3     |       |       |       |       |       |       |       |
| Patient placed in bed  | 2     |       |       |       |       |       |       |       |
| Out-patient area   | 1     |       |       |       |       |       |       |       |
| Patient had Surgery/Deep Sedation  |       |       |       |       |       |       |       |       |
| Within 24 hours  | 3     |       |       |       |       |       |       |       |
| Within 48 hours  | 2     |       |       |       |       |       |       |       |
| More than 48 hours/None  | 1     |       |       |       |       |       |       |       |
| Medication Usage   |       |       |       |       |       |       |       |       |
| Multiple usage of<br>Sedatives(Excluding ICU);<br>Hypnotics; Barbiturates;<br>Antidepressants;laxative;Diuretic<br>s; Narcotic | 3     |       |       |       |       |       |       |       |
| One of the medications listed<br>above   | 2     |       |       |       |       |       |       |       |
| Other Medications/None   | 1     |       |       |       |       |       |       |       |
| TOTAL SCORE  |       |       |       |       |       |       |       |       |
| Name & Signature   |       |       |       |       |       |       |       |       |

\*Low Fall Risk = 7-11 (Initiate low falls risk interventions)  
\*High Fall Risk = ≥ 12 (Initiate high falls risk interventions)